

Cydectin[®] Long Acting Injection for Sheep

Product Statement

For the treatment and control of roundworms, nasal bot and itchmite in sheep.
For protection against severe challenge by *Haemonchus contortus* (barber's pole worm) for up to 4 months

Composition

- Moxidectin 20g/L

What is Cydectin[®] Long Acting?

- Contains moxidectin – the most potent ML
- Not less than 91 days persistent activity against barber's pole worm and small brown stomach worm
- Up to 49 days persistent activity against black scour worm
- Ideal for highly contaminated pastures

Dosage

Cydectin[®] Long Acting Injection should be injected subcutaneously at a dose rate of 1mL/20kg liveweight.

Bodyweight (kg)	Dose Volume (mL)	Animals Treated
20*	1	500
21 - 30	1.5	333
31 - 40	2	250
41 - 50	2.5	200
51 - 60	3	166
61 - 70	3.5	142
71 - 80	4	125
81 - 90	4.5	111

DO NOT USE IN LAMBS LESS THAN 20KG.

Animals heavier than 90kg should be dosed at 0.5mL/10kg

Retreatment interval: must not be retreated for at least 91 days after administration

Withholding periods / ESI

- Meat: 91 days

- Milk: Do not use in female sheep which are producing, or may in the future produce, milk or milk products for human consumption
- ESI: 133 days

Pack Size

Liquid: 500mL

Storage


30°C (Room temperature) and protect from sunlight. Once opened contents should be used within 24hrs.

Manufacturer

Virbac Australia Pty Ltd

APVMA number

58532



For an obligation free quote on Cydectin® Long Acting Injection for Sheep, please visit the Animal Health Direct website at;

<http://www.animalhealthdirect.com.au/page/6/ContactUs.html>